



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b>  <b>A61K 31/44</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/55122</b>  <b>(43) International Publication Date:</b> 10 December 1998 (10.12.98)
<b>(21) International Application Number:</b> PCT/EP98/03478  <b>(22) International Filing Date:</b> 2 June 1998 (02.06.98)  <b>(30) Priority Data:</b> 9711683.4                      5 June 1997 (05.06.97)                      GB 9712851.6                      18 June 1997 (18.06.97)                      GB  <b>(71) Applicants (for all designated States except US):</b> SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19101 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PATEL, Jai [IN/US]; SmithKline Beecham Pharmaceuticals, 1250 South Collegeville Road, P.O. Box 5089, Collegeville, PA 19426-0989 (US). ROSS, Hamish [GB/US]; SmithKline Beecham Pharmaceuticals, 1250 South Collegeville Road, P.O. Box 5089, Collegeville, PA 19426-0989 (US). PRICE, Robin [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). GRANETT, Jeffrey, Roger [US/US]; SmithKline Beecham Pharmaceuticals, 1250		South Collegeville Road, P.O. Box 5089, Collegeville, PA 19426-0989 (US).  <b>(74) Agent:</b> RUTTER, Keith; SmithKline Beecham plc, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> COMPOSITION COMPRISING 5-[4-[2-(N-METHYL-N-2-PYRIDYL)AMINO]ETHOXY]BENZYL]THIAZOLIDINE-2,4-DIONE  <b>(57) Abstract</b>  A pharmaceutical composition comprising Compound (I), characterised in that the composition comprises 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form and optionally a pharmaceutically acceptable carrier therefor, the use of such a composition in medicine, processes for the preparation of such a composition and intermediate composition useful in such a process.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## COMPOSITION COMPRISING 5-[4-[2-(N-METHYL-N-2-PYRIDYL)AMINO)ETHOXY]BENZYL]THIAZOLIDINE-2,4-DIONE

This invention relates to a composition, in particular to a pharmaceutical composition, and to the use of such a composition in medicine, to processes for the preparation of such a composition and to a composition useful in such a process.

European Patent Application, Publication Number 0,306,228 relates to certain thiazolidinedione derivatives disclosed as having hypoglycaemic and hypolipidaemic activity. One particular thiazolidinedione disclosed in EP 0306228 is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (hereinafter 'Compound (I)'). International Patent Application, publication number WO94/05659 discloses certain salts of Compound (I), including the maleate salt at Example 1 thereof.

It is now surprisingly indicated that a discrete and particular daily dosage of Compound (I) provides an especially beneficial effect on glycaemic control and is therefore particularly useful for treatment of diabetes mellitus, especially Type II diabetes and conditions associated with diabetes mellitus.

We have also discovered a new and advantageous method for preparing pharmaceutical compositions, especially unit dosage compositions, containing Compound (I). The new method involves the preparation of a pre-administration concentrate of Compound (I) which thereafter is formulated into the required unit dose in an efficient and economical manner. The new process is particularly advantageous for the preparation of tablets of Compound (I).

Accordingly, in a first aspect the present invention provides a pharmaceutical composition, suitably in unit dosage form, comprising Compound (I), characterised in that the composition comprises 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form and optionally a pharmaceutically acceptable carrier therefor.

Suitable pharmaceutically acceptable forms of Compound (I) include pharmaceutically acceptable salted forms and pharmaceutically acceptable solvated forms, including pharmaceutically acceptable solvated forms of pharmaceutically acceptable salts.

Suitable compositions comprise 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 mg of Compound (I) in a pharmaceutically acceptable form.

Particular compositions comprise 2 to 4mg of Compound (I) in a pharmaceutically acceptable form.

Particular compositions comprise 4 to 8mg of Compound (I) in a pharmaceutically acceptable form.

Particular compositions comprise 8 to 12 mg of Compound (I) in a pharmaceutically acceptable form.

One composition comprises 2 mg of Compound (I) in a pharmaceutically acceptable form.

Preferred compositions comprise 4 mg of Compound (I) in a pharmaceutically acceptable form.

Preferred compositions comprise 8 mg of Compound (I) in a pharmaceutically acceptable form.

Suitable pharmaceutically acceptable salted forms of Compound (I) include those described in EP 0306228 and WO94/05659. A preferred pharmaceutically acceptable salt is a maleate.

Suitable pharmaceutically acceptable solvated forms of Compound (I) include those described in EP 0306228 and WO94/05659, in particular hydrates.

Compound (I) or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, may be prepared using known methods, for example those disclosed in EP 0306228 and WO94/05659. The disclosures of EP 0306228 and WO94/05659 are incorporated herein by reference.

Compound (I) may exist in one of several tautomeric forms, all of which are encompassed by the term 'Compound (I)' as individual tautomeric forms or as mixtures thereof.

Compound (I) contains a chiral carbon atom, and hence can exist in up to two stereoisomeric forms, the term Compound (I) encompasses all of these isomeric forms whether as individual isomers or as mixtures of isomers, including racemates.

When used herein the term 'conditions associated with diabetes' includes those conditions associated with the pre-diabetic state, conditions associated with diabetes mellitus itself and complications associated with diabetes mellitus.

When used herein the term 'conditions associated with the pre-diabetic state' includes conditions such as insulin resistance, including hereditary insulin resistance, impaired glucose tolerance and hyperinsulinaemia.

'Conditions associated with diabetes mellitus itself' include hyperglycaemia insulin resistance, including acquired insulin resistance and obesity. Further conditions associated with diabetes mellitus itself include hypertension, cardiovascular disease, especially atherosclerosis, certain eating disorders, in particular those requiring the regulation of appetite and food intake, such as disorders associated with under-eating, for example anorexia nervosa and disorders associated with over-eating, for example obesity and anorexia bulimia. Additional conditions associated with diabetes mellitus itself include polycystic ovarian syndrome and steroid induced insulin resistance and gestational diabetes.

'Complications associated with diabetes mellitus' includes renal disease, especially renal disease associated with Type II diabetes, including diabetic nephropathy, glomerulonephritis, glomerular sclerosis, nephrotic syndrome, hypertensive nephrosclerosis and end stage renal disease.

As used herein the term 'pharmaceutically acceptable' embraces both human and veterinary use: for example the term 'pharmaceutically acceptable' embraces a veterinarily acceptable compound.

As used herein the term concentrate with respect to Compound (I) in a pharmaceutically acceptable form means a proportionate amount of Compound (I) in a pharmaceutically acceptable form greater than that present in an administerable composition.

For the avoidance of doubt, when reference is made herein to scalar amounts, including mg amounts and % weight amounts, of 'Compound (I) in a pharmaceutically acceptable form', the scalar amount referred to is made in respect of Compound (I) *per se*: For example 2 mg of Compound (I) in the form of the maleate salt is that amount of maleate salt which contains 2 mg of Compound (I).

Diabetes mellitus is preferably Type II diabetes.

In a further aspect, the invention provides a process for preparing a pharmaceutical composition comprising 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form, and a pharmaceutically acceptable carrier therefor, which process comprises admixing 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form and the pharmaceutically acceptable carrier and optionally thereafter formulating the composition produced into an administerable form.

As indicated above the invention also provides a further process for preparing a pharmaceutical composition comprising Compound (I) in a pharmaceutically acceptable form which is particularly suitable for preparing a range of unit dosage forms of Compound (I). Accordingly, the invention further provides a process for preparing a pharmaceutical composition of Compound (I) in a pharmaceutically acceptable form and a pharmaceutically acceptable carrier, which process comprises:

- (i) preparing a first composition comprising Compound (I) in a pharmaceutically acceptable form and a first pharmaceutically acceptable carrier;
- (ii) admixing the first composition with a second pharmaceutically acceptable carrier to provide the required composition of Compound (I) and optionally thereafter formulating the composition produced into an administerable form.

A preferred administerable form of the pharmaceutical composition of Compound (I) is a unit dose composition.

Unless otherwise specified, suitable unit doses comprise up to 12 mg, such as 1 to 12 mg, of Compound (I) in a pharmaceutically acceptable form.

Other unit doses include those mentioned herein.

A key component of the last above mentioned process is the first composition. Accordingly, the present invention also provides a composition for use as a first composition in a process for preparing a unit dose of Compound (I) in a pharmaceutically acceptable form.

The invention also provides a composition comprising Compound (I) in a pharmaceutically acceptable form and optionally a pharmaceutically acceptable carrier, characterised in that the composition is a pharmaceutically acceptable, pre-administration composition.

A suitable pharmaceutically acceptable, pre-administration composition is a concentrate, preferably a granular concentrate, of Compound (I) in a pharmaceutically acceptable form. The granular concentrate is particularly well adapted to be diluted to provide a composition for administration, preferably a tablet.

In a further aspect the invention provides a composition comprising Compound (I) in a pharmaceutically acceptable form and a pharmaceutically acceptable carrier, characterised in that the composition is a concentrate of Compound (I) in a pharmaceutically acceptable form, adapted to be diluted so as to provide a composition for administration.

Suitably, the first composition, pre-administration composition or dilutable composition (hereinafter referred to for convenience as 'the first composition') contains up to 50% by weight, for example an amount in the range of from 2 to 50% by weight, of Compound (I) in a pharmaceutically acceptable form.

Favourably, the first composition contains an amount of Compound (I) in a pharmaceutically acceptable form in the range of from 5 to 20% by weight, in particular 5%, 10% or 15% by weight, for example 10% by weight.

The processes of the invention can provide pharmaceutical compositions of Compound (I) in any conventionally administerable form, including orally or parenterally administerable forms. They are particularly well adapted for preparing orally administrable forms, especially tablet forms of Compound (I) in a pharmaceutically acceptable form.

The first pharmaceutically acceptable carrier can comprise any conventional pharmaceutically acceptable carrier comprising conventional pharmaceutically acceptable excipients, including those disclosed in the reference texts mentioned below. However, as it is not essential that the first pharmaceutically acceptable carrier is in an administerable form, then it need not contain excipients solely associated with administration. For example the first pharmaceutically acceptable carrier need not contain a lubricant.

The second pharmaceutically acceptable carrier includes any conventional pharmaceutically acceptable carrier comprising any conventional pharmaceutically acceptable excipient, including disintegrants, diluents and lubricants, including those disclosed in the reference texts mentioned below.

One particular first composition comprises Compound (I) in a pharmaceutically acceptable form, a disintegrant, a binder and a diluent.

A suitable disintegrant is sodium starch glycollate.

A suitable binder is a methyl cellulose binder, such as hydroxypropyl methylcellulose 2910.

Suitable diluents include cellulose, for example a microcrystalline cellulose, and lactose monohydrate.

A suitable lubricant is magnesium stearate.

We have found that a particularly advantageous first composition contains Compound (I) in a pharmaceutically acceptable form, sodium starch glycollate, hydroxypropyl methylcellulose 2910, microcrystalline cellulose and lactose

monohydrate, especially when in a granular form. This granular form has been found to be particularly stable.

When the first composition contains about 10% by weight of Compound (I) in a pharmaceutically acceptable form, it is readily dilutable to give unit dose compositions comprising in the range of between 2 to 12 mg, especially 2 to 8 mg, 2 to 4mg, 4 to 8 mg and 8 to 12 mg of Compound (I) in a pharmaceutically acceptable form.

The preparation of the first composition is suitably carried using any conventional method appropriate to the nature of the said first composition, for example wet granulation methods provide the first composition in granular form.

Methods for formulating the compositions of the invention into administerable forms include conventional formulation methods as disclosed in the reference texts cited herein, including tableting methods.

The administerable compositions of the invention are preferably adapted for oral administration. However, they may also be adapted for other modes of administration, for example parenteral administration, sublingual or transdermal administration.

The administerable compositions may be in the form of tablets, capsules, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid preparations, such as oral or sterile parenteral solutions or suspensions.

In order to obtain consistency of administration it is preferred that a composition of the invention is in the form of a unit dose.

Unit dose presentation forms for oral administration may be tablets and capsules and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate; disintegrants, for example starch, polyvinylpyrrolidone, sodium starch glycollate or microcrystalline cellulose; or pharmaceutically acceptable wetting agents such as sodium lauryl sulphate.

Unless otherwise prescribed, compositions of the invention are preferably in unit dosage form in an amount appropriate for the relevant daily dosage, suitable unit dosages comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 mg of Compound (I) in a pharmaceutically acceptable form.

The solid compositions for example the oral compositions may be prepared by conventional methods of blending, filling or tableting. As required repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are of course conventional in the art. The tablets may be coated according to methods



well known in normal pharmaceutical practice, in particular with an aqueous film coating.

Liquid compositions, for example oral liquid compositions, may be in the form of emulsions, syrups, or elixirs, or they may be in a dry product form to be reconstituted with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel, hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid; and if desired conventional flavouring or colouring agents.

Parenteral compositions, including parenteral administration compositions for example unit dosage compositions, comprise the active compound and a sterile vehicle, and, depending upon the concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions for parenteral administration the composition of the invention may be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, a preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the active compound is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

Unless otherwise specified the compositions of the invention may contain from 0.1% to 99% by weight, preferably from 10-60% by weight, of the active material, depending upon the method of administration.

The composition may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

The compositions of the invention may be prepared and formulated according to conventional methods, such as those disclosed in standard reference texts, for example the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) and Harry's Cosmeticology (Leonard Hill Books).

The present invention also provides a pharmaceutical composition comprising 2 to 12 mg of Compound (I) and a pharmaceutically acceptable carrier therefor, for use as an active therapeutic substance.

In particular, the present invention provides a pharmaceutical composition comprising 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form, for use in the treatment of diabetes mellitus, especially Type II diabetes, and conditions associated with diabetes mellitus..

The composition of the invention may be administered from 1 to 6 times a day, but most preferably 1 or 2 times per day.

Accordingly, in a further aspect the invention provides a method for treatment of diabetes mellitus, especially Type II diabetes and conditions associated with diabetes mellitus, in a mammal such as a human, which method comprises administering per day 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form, to a mammal in need thereof.

Particularly, the method comprises the administration of 2 to 4 , 4 to 8 or 8 to 12 mg of Compound (I) in a pharmaceutically acceptable form.

Particular dosages are 2mg/day, 4mg/day, including 2mg twice per day, and 8 mg/day, including 4mg twice per day.

Particularly, the method comprises the administration of 2 to 4mg of Compound (I) in a pharmaceutically acceptable form.

Particularly, the method comprises the administration of 4 to 8mg of Compound (I) in a pharmaceutically acceptable form.

Particularly, the method comprises the administration of 8 to 12 mg of Compound (I) in a pharmaceutically acceptable form.

Preferably, the method comprises the administration of 2 mg of Compound (I) in a pharmaceutically acceptable form.

Preferably, the method comprises the administration of 4 mg of Compound (I) in a pharmaceutically acceptable form.

Preferably, the method comprises the administration of 8 mg of Compound (I) in a pharmaceutically acceptable form.

A range of 2 to 4mg includes a range of 2.1 to 4, 2.2 to 4, 2.3 to 4, 2.4 to 4, 2.5 to 4, 2.6 to 4, 2.7 to 4, 2.8 to 4, 2.9 to 4 or 3 to 4mg.

A range of 4 to 8mg includes a range of 4.1 to 8, 4.2 to 8, 4.3 to 8, 4.4 to 8, 4.5 to 8, 4.6 to 8, 4.7 to 8, 4.8 to 8, 4.9 to 8, 5 to 8, 6 to 8 or 7 to 8mg.

A range of 8 to 12 mg includes a range of 8.1 to 12, 8.2 to 12, 8.3 to 12, 8.4 to 12, 8.5 to 12, 8.6 to 12, 8.7 to 12, 8.8 to 12, 8.9 to 12, 9 to 12, 10 to 12 or 11 to 12mg.

No adverse toxicological effects have been established for the compositions or methods of the invention in the abovementioned dosage ranges.

The following examples illustrate the invention but do not limit it in any way.

**Example 1: Concentrate Preparation**

Approximately two thirds of the lactose monohydrate is passed through a suitable screen and blended with the milled maleate salt of Compound (I).

Sodium starch glycollate, hydroxypropyl methylcellulose, microcrystalline cellulose and the remaining lactose are passed through a suitable screen and added to the mixture. Blending is then continued. The resulting mixture is then wet granulated with purified water. The wet granules are then screened, dried on a fluid bed drier and the dried granules are passed through a further screen and finally homogenised.

**% COMPOSITION OF GRANULAR CONCENTRATE**

<b>Ingredient</b>	<b>Quantity (%)</b>
Milled Compound (I) as maleate salt	13.25 (pure maleate salt)
Sodium Starch Glycollate	5.00
Hydroxypropyl Methylcellulose 2910	5.00
Microcrystalline Cellulose	20.0
Lactose Monohydrate, regular grade	to 100
Purified water	*

\* Removed during processing.

Example 2: Formulation of the concentrate into tablets.

The granules from Example 1 are placed into a tumble blender. Approximately two thirds of the lactose is screened and added to the blender. The microcrystalline cellulose, sodium starch glycollate, magnesium stearate and remaining lactose are screened and added to the blender and the mixture blended together. The resulting mix is then compressed on a rotary tablet press to a target weight of 150mg for the 1, 2 and 4mg tablets and to a target weight of 300mg for the 8mg tablets.

The tablet cores are then transferred to a tablet coating machine, pre-warmed with warm air (approximately 65°C) and film coated until the tablet weight has increased by 2.0% to 3.5%.

Tablet Strength	Quantity (mg per Tablet)			
	1.0mg	2.0mg	4.0mg	8.0mg
<b>Active Ingredient:</b>				
Compound (I) maleate Concentrate granules	10.00	20.00	40.00	80.00
<b>Other Ingredients:</b>				
Sodium Starch Glycollate	6.96	6.46	5.46	10.92
Microcrystalline Cellulose	27.85	25.85	21.85	43.70
Lactose monohydrate	104.44	96.94	81.94	163.88
Magnesium Stearate	0.75	0.75	0.75	1.50
<b>Total Weight of Tablet Core</b>	150.0	150.0	150.0	300.0
Aqueous film coating material	4.5	4.5	4.5	9.0
<b>Total Weight of Film Coated Tablet</b>	154.5	154.5	154.5	309.0

## Claims:

1. A pharmaceutical composition comprising 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (hereinafter 'Compound (I)'), characterised in that the composition comprises 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form and optionally a pharmaceutically acceptable carrier therefor.
2. A composition according to claim 1, which comprises 2 to 4mg of Compound (I) in a pharmaceutically acceptable form.
3. A composition according to claim 1, which comprises 4 to 8mg of Compound (I) in a pharmaceutically acceptable form.
4. A composition according to claim 1, which comprises 8 to 12 mg of Compound (I) in a pharmaceutically acceptable form.
5. A composition according to claim 1, which comprises 2 mg of Compound (I) in a pharmaceutically acceptable form.
6. A composition according to claim 1, which comprises 4 mg of Compound (I) in a pharmaceutically acceptable form.
7. A composition according to claim 1, which comprises 8 mg of Compound (I) in a pharmaceutically acceptable form.
8. A composition according to any one of claims 1 to 7, which comprises the maleate salt of Compound (I)
9. A process for preparing a pharmaceutical composition comprising 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form, and a pharmaceutically acceptable carrier therefor, which process comprises admixing 2 to 12 mg of Compound (I) in a pharmaceutically acceptable form and the pharmaceutically acceptable carrier.

10. A process for preparing a pharmaceutical composition of Compound (I) in a pharmaceutically acceptable form and a pharmaceutically acceptable carrier, which process comprises:

- (i) preparing a first composition comprising Compound (I) in a pharmaceutically acceptable form and a first pharmaceutically acceptable carrier;
- (ii) admixing the first composition with a second pharmaceutically acceptable carrier to provide the required composition of Compound (I) and optionally thereafter formulating the composition produced into an administerable form.

11. A process according to claim 9 or claim 10, wherein the composition prepared is in unit dosage form.

12. A process according to claim 9 or claim 10, wherein the composition prepared is a tablet.

13. A composition for use as a first composition in a process according to claim 10, for preparing a unit dose of Compound (I) in a pharmaceutically acceptable form.

14. A composition comprising Compound (I) in a pharmaceutically acceptable form and optionally a pharmaceutically acceptable carrier, characterised in that the composition is a pharmaceutically acceptable, pre-administration composition.

15. A pre-administration composition according to claim 13, which is a concentrate of Compound (I) in a pharmaceutically acceptable form.

16. A composition comprising Compound (I) in a pharmaceutically acceptable form and a pharmaceutically acceptable carrier, characterised in that the composition is a concentrate of Compound (I) in a pharmaceutically acceptable form, adapted to be diluted so as to provide a composition for administration.

17. A composition according to any one of claims 13 to 16, which contains up to 50% by weight of Compound (I) in a pharmaceutically acceptable form.

18. A composition according to any one of claims 13 to 17, which contains an amount of Compound (I) in a pharmaceutically acceptable form in the range of from 5 to 20% by weight.

19. A composition according to any one of claims 13 to 18, which contains 5%, 10% or 15% by weight of Compound (I) in a pharmaceutically acceptable form.

20. A composition according to any one of claims 13 to 19, which contains Compound (I) in a pharmaceutically acceptable form, sodium starch glycollate, hydroxypropyl methylcellulose 2910, microcrystalline cellulose and lactose monohydrate.

21. A composition according to any one of claims 13 to 20, in granular form.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/03478

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	"ROSIGLITAZONE SMITHKLINE BEECHAM CLINICAL DATA" STN FILE SUPPLIER: DRUGNL: ACCESSION NUMBER 1391 & R&D FOCUS DRUG NEWS, 27 April 1998, XP002082989 see the whole document ---	1-4, 6-9
P, X	"THE PINK SHEET" DIALOG FILE SUPPLIER: FILE 187:F-D-C REPORTS; ACCESSION NUMBER 600160005, vol. 60, no. Issue 16, 20 April 1998, XP002082978 see the whole document --- -/--	1, 3, 4, 7-9

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "3" document member of the same patent family

Date of the actual completion of the international search

3 November 1998

Date of mailing of the international search report

17/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Hoff, P



## INTERNATIONAL SEARCH REPORT

In International Application No  
PCT/EP 98/03478

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 98 02159 A (POSTE GEORGE HENRY ;SMITH STEPHEN ALISTAIR (GB); SMITHKLINE BEECHA) 22 January 1998 see page 14, line 4 - line 10 see page 19, line 1 - line 16; claims; examples ---	1,4,8-16
X	WO 95 21608 A (SMITHKLINE BEECHAM PLC ;BUCKINGHAM ROBIN EDWIN (GB)) 17 August 1995 see page 4, line 28 - page 5, line 2 see page 7, line 29 - page 8, line 33; claims; examples ---	1,4,8-16
X	EP 0 796 618 A (SANKYO CO) 24 September 1997 see page 8, line 28 - line 30 see page 8, line 58 - page 9, line 42 ---	10-17
X	WO 97 05875 A (WARNER LAMBERT CO ;WHITCOMB RANDALL W (US)) 20 February 1997 see page 20, line 19 - line 33 see page 24, line 24 - page 27, line 8 ---	10-19,21
X	WO 97 18811 A (WARNER LAMBERT CO ;LOCKWOOD DEAN (US)) 29 May 1997 see page 18, line 15 - line 29 see page 21, line 16 - page 23, line 34 ---	10-19,21
Y	EP 0 306 228 A (BEECHAM GROUP PLC) 8 March 1989 cited in the application see page 9, line 9 - page 10, line 13; claims; example 30 ---	1-21
Y	WO 94 05659 A (SMITHKLINE BEECHAM PLC ;POOL COLIN RIPLEY (GB); ROMAN ROBIN SHERWO) 17 March 1994 cited in the application see the whole document ---	1-21
Y	"R&D FOCUS DRUG NEWS " DIALOG FILE SUPPLIER: FILE 445 R&D FOCUS; ACCESSION NUMBERS 16402 AND 14818, 5 May 1997 - 10 March 1997, XP002082979 see the whole document ---	1-21
	--- -/--	

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
A	<p>BERGER J ET AL: "THIAZOLIDINEDIONES PRODUCE A CONFORMATIONAL CHANGE IN PEROXISOMAL PROLIFERATOR-ACTIVATED RECEPTOR-GAMMA: BINDING AND ACTIVATION CORRELATE WITH ANTIDIABETIC ACTIONS IN DB/DB MICE" ENDOCRINOLOGY, vol. 137, no. 10, October 1996, pages 4189-4195, XP000613643 -----</p>	1-21

# INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/EP 98/03478

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9802159	A	22-01-1998	AU 3552697 A	09-02-1998
WO 9521608	A	17-08-1995	AP 553 A	06-11-1996
			AU 1578395 A	29-08-1995
			CA 2182986 A	17-08-1995
			CN 1145027 A	12-03-1997
			EP 0777469 A	11-06-1997
			HU 74382 A	30-12-1996
			JP 9512249 T	09-12-1997
			SG 47100 A	20-03-1998
			ZA 9501002 A	08-08-1996
EP 0796618	A	24-09-1997	AU 1628597 A	25-09-1997
			CA 2200126 A	18-09-1997
			CN 1170575 A	21-01-1998
			CZ 9700790 A	15-10-1997
			HU 9700598 A	28-09-1998
			JP 10147528 A	02-06-1998
			NO 971219 A	19-09-1997
			US 5753681 A	19-05-1998
WO 9705875	A	20-02-1997	AU 6641196 A	05-03-1997
			CA 2221241 A	20-02-1997
			EP 0851757 A	08-07-1998
			NO 980556 A	09-02-1998
WO 9718811	A	29-05-1997	AU 7604596 A	11-06-1997
EP 0306228	A	08-03-1989	AU 2173888 A	09-03-1989
			CA 1328452 A	12-04-1994
			DK 490288 A	05-03-1989
			EP 0842925 A	20-05-1998
			JP 10194970 A	28-07-1998
			JP 10194971 A	28-07-1998
			JP 1131169 A	24-05-1989
			JP 2614497 B	28-05-1997
			JP 9183771 A	15-07-1997
			JP 9183726 A	15-07-1997
			JP 9183772 A	15-07-1997
			PT 88410 A,B	31-07-1989

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0306228	A		US 5646169 A	08-07-1997
			US 5002953 A	26-03-1991
			US 5521201 A	28-05-1996
			US 5232925 A	03-08-1993
			US 5194443 A	16-03-1993
			US 5756525 A	26-05-1998
			US 5260445 A	09-11-1993
<hr/>				
WO 9405659	A	17-03-1994	AU 674880 B	16-01-1997
			AU 4973093 A	29-03-1994
			CA 2143849 A	17-03-1994
			CN 1101911 A	26-04-1995
			CN 1183275 A	03-06-1998
			CN 1183413 A	03-06-1998
			CN 1183276 A	03-06-1998
			CZ 9500565 A	15-11-1995
			EP 0658161 A	21-06-1995
			FI 951004 A	03-03-1995
			HU 72639 A	28-05-1996
			IL 106904 A	30-09-1997
			JP 8501095 T	06-02-1996
			MX 9305397 A	31-01-1995
			NO 950852 A	03-03-1995
			NO 974646 A	03-03-1995
			NZ 255505 A	22-08-1997
			PL 307812 A	26-06-1995
			SG 48302 A	17-04-1998
			SI 9300452 A	30-06-1994
			SK 27795 A	09-08-1995
			US 5741803 A	21-04-1998
			ZA 9306509 A	16-06-1994
<hr/>				